



# How to choose your Glucagon ELISA

What makes a glucagon assay great?

# How to choose your Glucagon ELISA

Glucagon is a secreted peptide hormone from alpha cells in the pancreas and it is an essential regulator of metabolic homeostasis. Glucagon has been established as an important hormone that orchestrates glucose, lipid, and amino acid metabolism, as well as energy intake and expenditure.

With Merckodia's experience developing premium immunoassays, the high specificity and sensitivity of the assays, makes it easier for researchers to get reliable results. Merckodia's gold-standard Glucagon ELISAs have been used by multiple researchers and clinicians worldwide, who appreciate the accuracy of the glucagon measurements.

**Inaccurate glucagon read-outs can result in problematic and negative implications on research and clinical decisions.**

## Superior sensitivity

Before using an assay, it is important to check the analytical sensitivity and ensure that the measurement range matches your expected sample levels. Some assays have a broader range than they can measure, so it is essential to confirm the accurate limits with the manufacturer.

Merckodia's Glucagon ELISAs have superior sensitivity, which enables you to measure physiologically suppressed levels of glucagon. This has been proven by multiple researchers, stating that only Merckodia's Glucagon ELISAs had optimal performance for measuring glucagon concentrations in clinical and animal samples (4, 5).

## High specificity

Any cross-reactivity to the proglucagon-derived gut peptides should be low and if present, well described since they contain the full glucagon sequence.

Glicentin circulates at much higher levels than glucagon, and therefore cross-reactivity can result in serious measurement errors. To avoid problems with your results, always confirm what concentrations were used in the manufacturer's cross-reactivity testing, since the level of cross-reactivity is usually positively correlated to the concentrations tested.

Merckodia's Glucagon ELISAs have no or very low cross-reactivity to glicentin, proglucagon, and oxyntomodulin.



### **Minimal matrix interference**

When analyzing samples, keep in mind that off-target molecules (like other proteins or lipids) could lead to altered or unexpected results.

Mercodia's Glucagon ELISAs contain a unique blocking solution to prevent or minimize matrix interferences, especially important for the animal glucagon assay.

### **Low sample volume**

Most commercially available methods require at least 50–100  $\mu\text{L}$  of plasma. Mercodia's Glucagon ELISAs were developed and optimized to require minimal sample volumes, with the assay for human samples requiring only 25  $\mu\text{L}$  per well and 10  $\mu\text{L}$  for the animal assay.

Low sample volumes enable scientists to measure glucagon along with other relevant hormones, and study temporal changes in glucagon in a variety of experimental models.

”

The validation of the Mercodia ELISA for measuring glucagon in human samples showed that the assay meets our established acceptability criteria.

**Kahn S.E. et al. 2021**

## References

1. Bak MJ, Albrechtsen NW, Pedersen J, *et al.* **Specificity and sensitivity of commercially available assays for glucagon and oxyntomodulin measurement in humans.** Eur J Endocrinol. 2014;170(4):529-538.
2. Campbell JE, Drucker DJ. **Islet  $\alpha$  cells and glucagon--critical regulators of energy homeostasis.** Nat Rev Endocrinol. 2015;11(6):329-338.
3. Howard JW, Kay RG, Tan T, Minnion J, Creaser CS. **Identification of plasma protease derived metabolites of glucagon and their formation under typical laboratory sample handling conditions.** Rapid Commun Mass Spectrom. 2015;29(2):171-181.
4. Wewer Albrechtsen NJ, Kuhre RE, Windeløv JA, *et al.* **Dynamics of glucagon secretion in mice and rats revealed using a validated sandwich ELISA for small sample volumes.** Am J Physiol Endocrinol Metab. 2016;311(2):E302-E309.
5. Kahn SE, Mather KJ, Arslanian SA, *et al.* **Hyperglucagonemia Does Not Explain the  $\beta$ -Cell Hyperresponsiveness and Insulin Resistance in Dysglycemic Youth Compared With Adults: Lessons From the RISE Study.** Diabetes Care. 2021;44(9):1961-1969.
6. Wewer Albrechtsen NJ, Kjeldsen SAS, Jensen NJ, *et al.* **On measurements of glucagon secretion in healthy, obese, and Roux-en-Y gastric bypass operated individuals using sandwich ELISA.** Scand J Clin Lab Invest. 2022;82(1):75-83.
7. Wewer Albrechtsen NJ, Holst JJ, Cherrington AD, *et al.* **100 years of glucagon and 100 more.** Diabetologia. 2023;66(8):1378-1394.

**mercodia.com**

Mercodia is a registered trademark of Mercodia AB.  
© Mercodia AB 2023. Mercodia AB, Sylveniusgatan 8A, SE 754 50 Uppsala, Sweden.

32-4255 v3 08/2023